

## 3-Colour Reagent anti-human CD4/CD3/CD8

### Premixed Antibody Combination

FITC conjugated anti human **CD4** (EDU-2; mouse IgG2a)  
PE conjugated anti human **CD3** (UCHT-1; mouse monoclonal IgG1/kappa)  
PerCP conjugated anti human **CD8** (UCHT-4; mouse IgG2a/kappa)

Cat-No: **21453006**

500 µl

**Specificity of CD4:** The antibody EDU-2 recognizes CD4 antigen, a 55 kDa transmembrane glycoprotein expressed on a subset of T lymphocytes ("helper" T-cells) and also on monocytes, tissue macrophages and granulocytes.

**Specificity of CD3:** The antibody UCHT1 recognizes the CD3 antigen of the TCR/CD3 complex on mature human T-cells. The UCHT1 antibody reacts with the epsilon chain of the CD3 complex.

**Specificity of CD8:** The UCHT-4 antibody recognises the CD8 alpha antigen on human cytotoxic/suppressor T-cells. On flow cytometry it stains 14-45% of human peripheral blood mononuclear cells. The antibody has been studied at the III. International Workshop on Human Leukocyte Differentiation Antigens.

CD8 is a TCR coreceptor molecule binding MHC class I. The CD8 antigen is expressed on human suppressor/cytotoxic T-cells, human NK-cell subset, cortical thymocyte subset and granulocytes.

**Form:** The anti-human CD4 antibody EDU-2 is conjugated with Fluorescein isothiocyanate (FITC). The anti-human CD3 antibody UCHT-1 is conjugated with R-Phycoerythrin (R-PE). The anti-human CD8 antibody UCHT-4 is conjugated with Peridin Chlorophyll.

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing BSA and 0.09% sodium azide (pH 7.2).

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

**Storage conditions:** Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.

**Application:** Flow Cytometry, 500µl = 25 tests

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Altenoyther Str. 10; 26169 Friesoythe; Germany

phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com

www.immunotools.com