

Recombinant human Soluble RANK Ligand (rh sRANKL / CD254)

Synonyms: Osteoprotegerin ligand (OPGL), Osteoclast differentiation factor (ODF), soluble Receptor Activator of Nuclear factor Kappa B Ligand, TNF-related activation-induced cytokine (TRANCE), Tumor necrosis factor ligand superfamily member 11 (TNFSF11)

Introduction: RANKL binds to tnfrsf11b/opg and to tnfrsf11a/rank. Osteoclast differentiation and activation factor. Augments the ability of dendritic cells to stimulate naive t-cell proliferation. May be an important regulator of interactions between t-cells and dendritic cells and may play a role in the regulation of the t-cell-dependent immune response. sRANKL may also play an important role in enhanced bone-resorption in humoral hypercalcemia of malignancy.

Description: Recombinant human RANKL (soluble) produced in E. coli is a single, non-glycosylated polypeptide chain containing 175 amino acids and having a molecular mass of 19.7 kDa.

Source: *Escherichia Coli*.

Physical Appearance: Sterile filtered white lyophilized (freeze-dried) powder.

Formulation: Lyophilized from a concentrated (1mg/ml) solution containing 10mM sodium phosphate, pH-7.5. The aliquots of 1µg and 2µg contain Trehalose 5% (w/vol) for better recovery.

Solubility: It is recommended to reconstitute the lyophilized rh sRANKL in sterile H₂O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability: Lyophilized rh sRANKL although stable at room temperature for 3 weeks, should be stored desiccated below -18° C. Upon reconstitution rh sRANKL should be stored at 4° C between 2-7 days and for future use below -18°. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Amino acid sequence: EKAMVDGSW LDIAKRSKLE AQPFAHLTIN ATDIPSGSHK VSLSSWYHDR
GWAKISNMTF SNGKLIVNQD GFYYLYANIC FRHHETSGDL ATEYLQLMVY VTKTSIKIPS SHTLMKGGST
KYWSGNSEFH FYSINVGGF KLRSGEEISI EVSNPSLLDP DQDATYFGAF KVRDID.

Biological Activity: The activity of RAW-Blue was measured to be 38.8 ng/ml, corresponding to a specific activity of 2.6×10^4 units/mg.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

<i>small</i>	2 µg	Cat.N°	11343452
<i>medium</i>	10 µg	Cat.N°	11343453
<i>large</i>	50 µg	Cat.N°	11343455

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone: +49-(0)4491-400997, fax: +49-(0)4491-400998, info@immunotools.com
www.immunotools.com