

Recombinant Mouse Adiponectin HEK derived (rm Acrp30 HEK)

Synonyms: APM-1.

Introduction: Adiponectin is the product of the apM1 gene which is physiologically active and specifically and highly expressed in adipose cells (Adipokine). The protein belongs to the soluble defense collagen super family; it has a collagen-like domain structurally homologous with collagen VIII and X and complement factor C1q-like globular domain. Adiponectin is produced and secreted exclusively by adipocytes, and is a relatively abundant plasma protein, accounting for up to 0.05% of total serum protein. Adiponectin forms homotrimers, which are the building blocks for higher order complexes found circulating in serum. It is capable of decreasing hyperglycemia and reversing insulin resistance.

Description: Mouse recombinant Adiponectin produced in HEK293 is a single polypeptide chain containing 244 amino acids. Adiponectin is fused with FLAG-tag.

Source: HEK293(Human Embryonic Kidney cell line)

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation: Lyophilized from a sterile filtered 0.4µm solution in PBS buffer
The aliquots of 1µg and 2µg contain Trehalose 5% (w/vol) for better recovery

Solubility: Add 0.2 ml of deionized water and let the lyophilized pellet dissolve completely.

Stability: For long term store lyophilized Adiponectin at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C. The lyophilized protein remains stable for 2 4 months when stored at -20°C.

Purity: Greater than 98.0% as determined by HPLC and SDS-PAGE.

Amino Acid Sequence: ETTTQGGPVL LPLPKGACTG WMAGIPGHPG HNGAPGRDGR DGTPGEKGEK
GDPGLIGPKG DIGETGVPGA EGPRGFPGIQ GRKGEPGEGA YVYRSAFSVG LETYVTIPNM PIRFTKIFYN
QQNHYDGSTG KFHCNIPGLY YFAYHIVYMK DVKVSFLFKK KAMLFTYDQY QENNVDQASG SVLLHLEVG
D QVWLQVYGE ERNGLYADND NDSTFTGFL YHDTNDYKDDDDK.

Biological Activity: Full-length mouse adiponectin activates AMP-activated protein kinase in hepatocyte and activates AMPK in HepG2 human hepatocytes at a concentration of 1 µg/ml.

Mouse adiponectin mammalian cell derived inhibits glucose production as shown by in-vitro gluconeogenesis assay in primary rat hepatocytes.

Application: ELISA, Western blotting, cell culture and/or animal studies.

This material is offered for research only. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

<i>small</i>	2 µg	Cat.N°	12344462
<i>medium</i>	10 µg	Cat.N°	12344463

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com