

Recombinant Mouse Eotaxin-2 (CCL24)

Synonyms: C-C motif chemokine 24 (CCL24), Small-inducible cytokine A24, Myeloid progenitor inhibitory factor 2 (MPIF-2), CK-beta-6 (Ckb-6), Eosinophil chemotactic protein 2

Introduction: Eotaxin-2 is a novel CC chemokine produced by activated monocytes and T lymphocytes. Eotaxin-2 selectively chemoattracts cells expressing CCR3 including eosinophils, basophils, Th2 T cells, mast cells and certain subsets of dendritic cells. Furthermore Eotaxin-2 inhibits the proliferation of multipotential hematopoietic progenitor cells. The mature protein which includes C-terminal truncation contains 78 amino acids (92 amino acids for the mouse homolog, without C-terminal truncation).

Eotaxin-2 functions as a chemotactic chemokine for resting T-lymphocytes and eosinophils. It has lower chemotactic activity for neutrophils but none for monocytes and activated lymphocytes. Eotaxin-2 is a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line and binds to CCR3.

Description: Recombinant mouse Eotaxin-2 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 93 amino acids and having a molecular mass of 10.3 kDa. The Eotaxin-2 is purified by proprietary chromatographic techniques.

Source: *Escherichia Coli*

Physical Appearance: Sterile filtered white lyophilized (freeze-dried) powder.

Formulation: Lyophilized from a concentrated solution containing 20mM PBS pH-7.4 and 0.15M NaCl. The sample size of 1µg contains Trehalose 5% (w/vol) for better recovery

Stability: Lyophilized Eotaxin-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL24 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

Purity: Greater than 97.0% as determined by:

- (a) Analysis by RP-HPLC.
- (c) Analysis by SDS-PAGE.

Amino Acid Sequence: VTIPSSCCTS FISKKIPENR VVSYQLANGS ICPKAGVIFI TKKGHKICTD PKLLWVQRHI QKLDKKNQP SKGAKAVRTK FAVQRRRGNS TEV.

Biological Activity: The activity is determined by its ability to chemoattract murine lymphocytes using a concentration range of 10ng/ml -100ng/ml.

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

<i>small</i>	5 µg	Cat.N°	12344170
<i>medium</i>	20 µg	Cat.N°	12344174

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
 phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com