

Recombinant Murine Interleukin-10 (rm IL-10)

Synonyms : Cytokine synthesis inhibitory factor (CSIF), B-TCGF, TGIF, IL10A

Introduction : IL10 is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down- regulates the expression of Th1 cytokines, MHC class II Ags and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation and antibody production. This cytokine can block NF-kappa B activity and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract.

Description: Recombinant murine IL-10 produced in *E.Coli* is a single, glycosylated polypeptide chain containing 161 amino acids and having a molecular mass of 18.8 kDa. The rm IL-10 is purified by proprietary chromatographic techniques.

Source: *Escherichia Coli*.

Physical Appearance: Sterile filtered white lyophilized (freeze-dried) powder.

Formulation: Lyophilized from a 0.22 µm filtered solution in TRIS, pH 8.0 + 250 mM NaCl. The aliquots of 1µg and 2µg contain Trehalose 5% (w/vol) for better recovery

Solubility: It is recommended to reconstitute the lyophilized rm IL-10 in sterile H₂O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Stability: Lyophilized rm IL-10 although stable at room temperature for 3 weeks, should be stored desiccated below -18° C. Upon reconstitution rm IL-10 should be stored at 4° C between 2-7 days and for future use below -18° C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze-thaw cycles.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Amino Acid Sequence: MSRGQYSRED NNCTHFPVGQ SHMLLELRTA FSQVKTFQKTDKQLDNILLT DSLMQDFKGY LGCQALSEMI QFYLVEVMPQ AEKHGPEIKE HLNSLGQKLL TLRMLRRCH RFLPCENKSK AVEQVKSDFN KLEDQGVYKA MNEFDIFINC IEAYMMIKMK S

Biological Activity: The ED₅₀ as determined by dose dependant proliferation of a murine MC/9 cell line was found to be < 2 ng/ml, corresponding to a specific activity of 5x10⁵ IU/mg.

Endotoxicity: The endotoxin level is less than 0.1 ng per µg (1EU/µg) determined by LAL method

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

<i>small</i>	2 µg	Cat.N°	12340102
<i>medium</i>	10 µg	Cat.N°	12340103
<i>large</i>	50 µg	Cat.N°	12340105
<i>x-large</i>	250 µg	Cat.N°	12340107
<i>xx-large</i>	1000 µg	Cat.N°	12340108

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
 phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com