

Recombinant Mouse Interleukin-21 (rm IL-21)

Synonymes: none

Introduction: IL-21 is produced by CD4+ T cells in response to antigenic stimulation. Its action enhances antigen-specific responses of immune cells. The biological effects of IL-21 include induction of differentiation of T-cells-stimulated B-cells into plasma cells and memory B-cells, stimulation (in conjunction) with IL-4 of IgG production and induction of apoptotic effects in native B-cells and stimulated B-cells in the absence of T-cell signaling. Additionally IL-21 promotes the anti-tumor activity of CD8+ T-cells and NK cells. IL-21 exerts its effect through binding to a specific type I cytokine receptor, IL-21R, which also contains the gamma chain and is found in other cytokine receptors including IL-2, IL-4, IL-7, IL-9 and IL-15. The IL-21/IL-21R interaction triggers a cascade of events which includes activation of the tyrosine kinases JAK1 and JAK3, followed by activation of the transcription factors STAT1 and STAT3.

Description: Recombinant murine IL-21 produced in *E.Coli* is a single, non-glycosylated polypeptide chain containing 130 amino acids and having a total molecular mass of 15 kDa. The rm IL-21 is purified by proprietary chromatographic techniques.

Source: *Escherichia Coli*.

Physical Appearance: Sterile filtered white lyophilized (freeze-dried) powder.

Formulation: Lyophilized from a 0.22 µm filtered solution in 25 mM sodium phosphate pH 7.3 + 200mM NaCl. The aliquots of 1µg and 2µg contain Trehalose 5% (w/vol) for better recovery.

Solubility: It is recommended to reconstitute the lyophilized rm IL-21 in sterile H₂O not less than 100µg/ml which can then be further diluted to other aqueous solutions.

Stability: Lyophilized rm IL-21 although stable at room temperature for 3 weeks, should be stored desiccated below -18° C. Upon reconstitution rm IL-21 should be stored at 4° C between 2-7 days and for future use below -18° C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Endotoxicity: The endotoxin level is less than 0.1 ng per µg (1EU/µg) determined by LAL method.

Amino Acid Sequence: MHKSSPQGPD RLLIRLRHLLI DIVEQLKIYE NDLPELLSA PQDVKGHCEH AAFACFQKAK LKPSNPGNNK TFIIDLVAQL RRRLPARRGG KKQKHIKCP SCDSYEKRTP KEFLERLKWLLQKMIHQHLS

Biological Activity: Tested by dose dependent stimulation of IL-21 induced reporter gene activation in IL-21 receptor transfected cells the specific activity of rm IL-21 is > 0.5x10⁵ IU/mg

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

<i>small</i>	2 µg	Cat.N°	12340212
<i>medium</i>	10 µg	Cat.N°	12340213
<i>large</i>	50 µg	Cat.N°	12340215
<i>x-large</i>	250 µg	Cat.N°	12340217
<i>xx-large</i>	1000 µg	Cat.N°	12340218

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
 phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com