

Recombinant Murine Stromal Cell-Derived Factor-1 alpha (CXCL12 α / rm SDF-1 alpha)

Synonyms: Pre-B cell growth-stimulating factor, PBSF, hIRH, TPAR1, SCYB12, TLSF-a.

Introduction: SDF-1 is produced in two forms, SDF-1 α /CXCL12a and SDF-1 β /CXCL12b, by alternate splicing of the same gene. The SDF-1 proteins belong to the group of CXC chemokines whose initial pair of cysteines are separated by one intervening amino acid. SDF-1 is strongly chemotactic for lymphocytes and has been implicated as an important cell co-ordinator during development. During embryogenesis it directs the migration of hematopoietic cells from foetal liver to bone marrow. Mice which were knocked-out for SDF-1 gene were lethal before the birth or within just 1 hour of life. As another role, SDF-1 alters also the electrophysiology of neurons. SDF-1 was shown to be expressed in many tissues in mice (including brain, thymus, heart, lung, liver, kidney, spleen and bone marrow). The receptor for this chemokine is CXCR4. This SDF-1-CXCR4 interaction used to be considered exclusive (unlike for other chemokines and their receptors), but recently it was suggested that SDF-1 is also bound by CXCR7 receptor. In human and mouse both SDF-1 and CXCR4 show high identity of sequence: 99% and 90%, respectively.

Description: Recombinant murine SDF-1alpha produced in *E. Coli* is a non-glycosylated, polypeptide chain containing 68 amino acids and having a molecular mass of 7.9 kDa. The rm SDF-1alpha is purified by proprietary chromatographic techniques.

Source: *Escherichia Coli*.

Physical Appearance: Sterile filtered white lyophilized (freeze-dried) powder.

Formulation: lyophilized from a 0.22 μ m filtered solution in 25 mM Sodiumphosphate, 100 mM NaCl pH 7.4. The aliquots of 1 μ g and 2 μ g contain Trehalose 5% (w/vol) for better recovery.

Solubility: It is recommended to reconstitute the lyophilized rm SDF-1alpha in sterile H₂O not less than 100 μ g/ml, which can then be further diluted to other aqueous solutions.

Stability: Lyophilized rm SDF-1alpha although stable at room temperature for 3 weeks, should be stored desiccated below -18° C. Upon reconstitution rm SDF-1alpha should be stored at 4° C between 2-7 days and for future use below -18° C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Purity: Greater than 95.0% as determined by SDS-PAGE

Amino acid Sequence: KPVLSYRCP CRFFESHAR ANVKHLKILN TPNCALQIVA RLKNNNRQVC
IDPKLKWIQE YLEKALNK.

Endotoxicity: The endotoxin level is less than 1 EU / μ g determined by LAL method

Biological Activity: Determined by chemotaxis assay with human PBMCs the ED50 has been 80 – 100 ng/ml.

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

<i>small</i>	2 μ g	Cat.N°	12343362
<i>medium</i>	10 μ g	Cat.N°	12343363
<i>large</i>	50 μ g	Cat.N°	12343365
<i>x-large</i>	250 μ g	Cat.N°	12343367
<i>x-large</i>	1000 μ g	Cat.N°	12343368

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone: +49-(0)4491-400997, fax: +49-(0)4491-400998, info@immunotools.com
www.immunotools.com