

anti-mouse Gr-1 APC-conjugated

APC- conjugated monoclonal antibody RB6-8C5 to mouse Gr-1

Cat-No: **22155246**

500 µl

Clone: RB6-8C5

Specificity: The anti-mouse Gr-1 monoclonal antibody reacts with the myeloid differentiation antigen GR-1. This 25-30 kDa cell surface antigen is expressed on myeloid cells but not lymphoid or erythroid cells. The expression of the Gr-1 antigen increases with granulocyte maturation as shown by the distinct populations of bone-marrow cells this monoclonal antibody labels: negative, low positive and high positive. Expression is transient on cells of monocytic lineage. The antibody is a useful antibody for studies of myeloid differentiation stages and their regulations by cytokines.

Isotype subclass: Rat IgG2b

Form: The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 1 % BSA and 0.09 % sodium azide (pH 7.2)

Expiration date: The reagent is stable until the expiry date stated on the vial label

Storage conditions: Store at 4 °C. Avoid prolonged exposure to light.

Application: Flow Cytometry

References:

- 1.) Spangrude, G.J., et al. Purification and characterization of mouse hematopoietic stem cells. *Science* 241:58-62, 1991.
- 2.) Muller, C. E. et al. Isolation of two early B lymphocyte progenitors from mouse marrow: a committed pre-pre-B cell and a clonogenic thy-1 lo hematopoietic stem cell. *Cell* 44:653-662.
- 3.) H. Kjetil et al. Characterization and regulation of RB6-8C5 antigen expression on murine bone marrow cells. 1991. *J. Immunol.* 147: 22-28.
- 4.) Brummer, Elmer et al. Immunological activation of polymorphonuclear neutrophils for fungal killing: Studies with murine cells and blastomyces dermatitidis in vitro. *J. Leuko. Bio* 36:505-520, 1984.
- 5.) Jutila, M.A. et al. Ly-6C is a monocyte/macrophage and endothelial cell differentiation antigen regulated by interferon-gamma. *J. Immunol.* 18: 1819-1826, 1988.
- 6.) Whiteland, J. L. et al (1995). Immunohistochemical detection of T cell subsets and other leukocytes in paraffin embedded rat and mouse tissues with monoclonal antibodies, *J.Histochem. Cytochem.* 43:313-320.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Spreestraße 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com