

anti-human CD42b APC-conjugated

APC - conjugated monoclonal antibody HIP1 to human CD42b

Cat-No: **21270426**

500 µl

Clone: HIP1

Specificity: The antibody HIP1 recognizes a glycoprotein, heterodimer (GPIb α) which contains an α chain (CD42b, 145 kDa) and a β chain (CD42c, 22 kDa) linked by a disulfid bond forming a noncovalent complex with CD42a (GPIX) and d. This antigen is expressed on platelets and megakaryocytes. The GPIb/IX complex (CD42a-d) serves as a receptor for von Willebrand Factor (vWF) and thrombin, and mediates adhesion of platelets to subendothelium of damaged vascular walls. The absence of the CD42 complex leads to the Bernard-Soulier Syndrome (BSS). HIP1 McAv inhibits the ristocetin-dependent binding of vWF to platelets, ristocetin-induced platelets agglutination, and partially inhibits collagen-induced aggregation.

Isotype subclass: Mouse IgG1

Form: The purified antibody is conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 1% BSA and 0.09% sodium azide (pH 7.2)

Expiration date: The reagent is stable until the expiry date stated on the vial label

Storage conditions: Store at 4 °C. Avoid prolonged exposure to light.

Application: Flow Cytometry

References: Leucocyte Typing IV. Knapp W et al. (Eds.), Oxford University Press (1989).

Background: **CD42b** (GPIb α) composes together with GPIb β , GPIX and GPV the GPIb-IX-V receptor complex critical in the process of platelet-rich thrombus formation by tethering the platelet to a thrombogenic surface. CD42b binds to von Willebrand factor (VWF) exposed at a site of vascular injury, as well as to thrombin, coagulation factors XI and XII, high molecular weight kininogen, TSP-1, integrin Mac-1 and P-selectin. The extracellular domain of CD42b by its interactions also contributes to metastasis.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com