

anti-human CD4 Biotin -conjugated

Biotin-conjugated monoclonal antibody to human CD4

Cat-No: **21270042**

50 µg in 100 µl

Clone: MEM-241

Specificity: The antibody MEM-241 recognizes CD4 antigen, a 55 kDa transmembrane glycoprotein expressed on a subset of T-cells (helper T-cells) and also on monocytes, tissue macrophages and granulocytes. HCDM (former HLDA VIII) Meeting, May 2006, Québec, Canada; WS Code M241

Isotype subclass: Mouse IgG1

Form: The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 0.09 % sodium azide (pH 7.2).

Storage conditions: Store at 4°C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.

Application: Flow Cytometry, Western Blotting

Expiration date: The reagent is stable until the expiry date stated on the vial label.

References:

- * Millan J, Cerny J, Horejsi V, Alonso MA: CD4 segregates into specific detergent-resistant T-cell membrane microdomains. Tissue Antigens. 1999 Jan;53(1):33-40.
- * Foti M, Phelouzat MA, Holm A, Rasmussen BJ, Carpentier JL: p56Lck anchors CD4 to distinct microdomains on microvilli. Proc Natl Acad Sci U S A. 2002 Feb 19;99(4):2008-13.

Background: CD4 is a single chain transmembrane glycoprotein and belongs to immunoglobulin supergene family. In extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). Transmembrane region forms 25 aa, cytoplasmic tail consists of 38 aa. Domains 1,2 and 4 are stabilized by disulfide bonds. The intracellular domain of CD4 is associated with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains.

Extracellular ligands: MHC class II molecules (binds to CDR2-like region in CD4 domain 1); HIV envelope protein gp120 (binds to CDR2-like region in CD4 domain 1); IL-16 (binds to CD4 domain 3), Human seminal plasma glycoprotein gp17 (binds to CD4 domain 1), L-selectin - Intracellular ligands: p56Lck

CD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection (human immunodeficiency virus; CD4 is primary receptor for HIV-1 surface glycoprotein gp120). CD4 regulates T-cell activation, T/B-cell adhesion, T-cell differentiation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone: +49-(0)4491-400997, fax: +49-(0)4491-400998, info@immunotools.com
www.immunotools.com