## anti-human CD19 Biotin -conjugated

Biotin-conjugated monoclonal antibody to human CD19

Cat-No: **21270192** 50 μg in 100 μl

Clone: LT19

**Specificity:** The antibody LT19 reacts with CD19 (B4), a 95 kDa type I transmembrane glycoprotein (immunoglobulin superfamily) expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.

Isotype subclass: Mouse IgG1

Form: The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free

of unconjugated biotin.

Physical state: Liquid

**Buffer/Additives/Preservative:** PBS containing 0.09% sodium azide (pH 7.2)

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

**Storage conditions:** Store at 4 °C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.

Application: Flow Cytometry, Immunoprecipitation

**Background: CD19** is a transmembrane glycoprotein of Ig superfamily expressed by B cells from the time of heavy chain rearrangement until plasma cell differentiation. It forms a tetrameric complex with CD21 (complement receptor type 2), CD81 (TAPA-1) and Leu13. Together with BCR (B cell antigen receptor), this complex signals to decrease B cell treshold for activation by the antigen. Besides being signal-amplifying coreceptor for BCR, CD19 can also signal independently of BCR coligation and it turns out to be a central regulatory component upon which multiple signaling pathways converge. Mutation of the CD19 gene results in hypogammaglobulinemia, whereas CD19 overexpression causes B cell hyperactivity.

## References:

- \*) Elias F,Flo J,Lopez RA,Zorzopulos J,Montaner A,Rodriguez JM:Strong cytosine-guanosine-independent immunostimulation in humans and other primates by synthetic oligodeoxynucleotides with PyNTTTTGT motifs. J Immunol.2003 Oct 1;171(7):3697-704.
- \*) Lin CW,Liu TY,Chen SU,Wang KT,Medeiros LJ,Hsu SM:CD94 1A transcripts characterize lymphoblastic lymphoma/leukemia of immature natural killer cell origin with destinct clinical features. Blood.2005 Nov 15;106(10):3567-74. Epub 2005 Jul 26.

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for <u>research use only</u>. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.