

## anti-rat CD4 FITC-conjugated

**FITC-** conjugated monoclonal antibody W3/25 to rat CD4

Cat-No: **23150043**

500 µl

**Clone:** W3/25

**Specificity:** This anti-rat T helper cell monoclonal antibody recognizes a determinant on the majority of thymocytes (90-95%), a subset of peripheral T cells and peritoneal macrophages. The antigen recognized by this antibody is a surface glycoprotein of Mr 48 000 – 52 000 and is the homologue of the human CD4 and the mouse L3/T4 antigen. This antibody labels the rat T helper subset, which mediates the helper activity for B and T cells, grafts vs. host (GVH) reactivity and produces IL-2 in the mixed lymphocyte reaction (MLR). Addition of the antibody to the MLR, inhibited proliferation and blocks the production of IL-2. T cells which mediate cytotoxicity and suppressor functions are not labelled. This clone is invaluable for separating T cell subsets for functional studies. It has been studying the role of T lymphocytes in graft rejection and in studying the subsets of T cells in the rat which mediate graft vs. host disease. This antibody is one of three antibodies which labels T lymphocyte populations in the rat. These clones include W3/13, which labels all T cells, as well as MRC OX-8 and W3/25 which label non-overlapping T cell subpopulations. These monoclonal antibodies used in concert are being employed extensively to investigate cellular aspects of the immune response in rats and prove to be useful as markers of functionally distinct subpopulations of lymphocytes.

**Isotype subclass:** Mouse IgG1

**Form:** The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing 1 % BSA and 0.09 % sodium azide (pH 7.2).

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

**Storage conditions:** Store at 4 °C. Avoid prolonged exposure to light.

**Application:** Flow Cytometry

### References:

1. Barclay, A.N. (1981) Immunology, 42, 593-600.
2. Cantrell, D.A., Robins, R.A. and R.W. Baldwin. (1982) Cell Immunol. 70, 367-372.
3. Mason, D.W., Pugh, C.W. and M. Webb. (1981) Immunology, 44, 75-87.
4. Dallman, M.J., Mason, D.W. and M. Webb. (1982) Eur. J. Immunol. 12, 511-518
5. Mason, D.W. (1981) Transplantation, 32, 222-226.
6. White, R.A.H., Mason, D.W., Williams, A.F., Galfre, G. and C. Milstein. (1978) J. Exp. Med. 148, 644-673
7. Jefferies, W.A., Green, J.R. and A.F. Williams. (1985) J. Exp. Med. 162, 117-127.
8. Whiteland, J.L. et al (1995). J.Histochem. Cytochem. 43: 313-320

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink, and animal feedingstuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany  
phone: +49-(0)4491-400997, fax: +49-(0)4491-400998, [info@immunotools.com](mailto:info@immunotools.com)  
[www.immunotools.com](http://www.immunotools.com)