

## anti-human CD50 FITC-conjugated

**FITC** - conjugated monoclonal antibody MEM-171 to human CD50

Cat-No: **21270503**

500 µl

**Clone:**MEM-171

**Specificity:** The antibody MEM-171 recognizes an epitope in the D2 domain of CD50 (ICAM-3), a 120-130 kDa type I membrane protein (immunoglobulin supergene family) expressed on leukocytes, endothelial cells and Langerhans cells; it is negative on platelets and erythrocytes.

**HLDA VI; WS Code BP 614**

**HLDA VI; WS Code NL N-L022**

**Isotype subclass:** Mouse IgG1

**Form:** Purified from ascites by protein-A affinity chromatography. Purified IgG, FITC conjugated

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing 1% BSA and 0.09% sodium azide (pH 7.2)

**Expiration date:** The reagent is stable until the expiry date stated on the vial label

**Storage conditions:** Store at 4 °C. Avoid prolonged exposure to light.

**Application:** Flow Cytometry

### References:

- \*Campanero MR and others: 1993 Nov;123(4):1007-16.
- \*Hernandez-Caselles T and others: 1993 Nov;23(11):2799-806.
- \*Arroyo AG and others: 1994 Sep;126(5):1277-86.
- \*Bogoevska V and others: 2007 Mar;17(3):324-33.).

**Background:** **CD50** (intracellular adhesion molecule 3, ICAM-3) is a transmembrane glycoprotein expressed by leukocytes, that serves as a counter-receptor for the lymphocyte function-associated antigen (LFA)-1 integrin. Besides functioning as an adhesive molecule that mediates e.g. the contact between T cells and antigen presenting cells, ICAM-3 regulates affinity of LFA-1 for ICAM-1 and induces T cell activation and proliferation. ICAM-3 plays an essential role in the initiation of the immune response both on T cells and antigen presenting cells and interacts also with CD209 (dendritic cell-specific ICAM-3-grabbing nonintegrin, DC-SIGN), a C-type lectin of dendritic cells and macrophages; this process is involved in dialogue between dendritic cells and granulocytes.

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany  
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com  
www.immunotools.com