

# anti-human CD53 FITC-conjugated

## FITC – conjugated monoclonal antibody MEM-53 to CD53 (Human)

Cat-No: **21270533**

500 µl

**Clone:** MEM-53

**Specificity:** The antibody MEM-53 reacts with CD53, a 32-40 kDa tetraspanin family glycoprotein exclusively expressed on leukocytes; it is not present on platelets, red blood cells and non-hematopoietic cells. The antibody MEM-53 reacts also with deglycosylated molecule (molecular weight of the antigen is reduced by 15 kDa using endoglycosidase F).

**HLDA IV; WS Code NL 59, HLDA V; WS Code B CD53.5 ,HLDA V; WS Code BP BP287  
HLDA V; WS Code T T-096, HLDA V; WS Code X XB004**

**Isotype subclass:** Mouse IgG1

**Form:** The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing 1% BSA and 0.09% sodium azide (pH 7.2)

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

**Storage conditions:** Store at 4 °C. Avoid prolonged exposure to light.

**Application:** Flow Cytometry

- References:**
- \*Angelisova P and others: 1990;32(4):281-5.
  - \*Olweus J and othes: 1993 Jul 15;151(2):707-16.
  - \*Mollinedo F: 1998 Jun;63(6):699-706.
  - \*Puls KL: 2002 Mar;14(3):249-58.
  - \*Kim TR.: 2004 Feb 29;17(1):125-31.
  - \*Yunta M, Lazo PA: 2003 Feb 27;22(8):1219-24.
  - \*Yunta M and others: 2003 Feb;63(2):534-42.

**Background:** **CD53** is a tetraspanin family transmembrane glycoprotein expressed in the lymphoid-myeloid lineage. This molecule has been reported to form complexes with other leukocyte surface proteins such as CD2, CD19, CD21, MHC II, VLA-4 or tetraspanins CD37, CD81 and CD82, thus probably modulating various signaling processes. CD53 is involved in radioresistance of tumour cells and its triggering has anti-apoptotic effect. In thymus, CD53 is up-regulated in response to positive selection signals during T cell development, and is strongly expressed upon macrophage exposure to bacterial lipopolysaccharide, whereas stimulation of neutrophils results in down-regulation of CD53 expression.

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany  
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com  
www.immunotools.com