## anti-human CD9 FITC-conjugated

FITC - conjugated monoclonal antibody HI9a to human CD9

Cat-No: **21810093** 500 μl

Clone: HI9a

**Specificity:** The antibody HI9a recognizes an epitope on second extracellular domain (EC2) of CD9 antigen, a 24 kDa transmembrane protein expressed on platelets, monocytes, pre-B lymphocytes, granulocytes and activated T lymphocytes.

Isotype subclass: Mouse IgG1

**Form:** The purified antibody is conjugated with Fluoresceinisothiocyanate (FITC) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 1% BSA and 0.09% sodium azide (pH 7.2)

**Expiration date:** The reagent is stable until the expiry date stated on the vial label

Storage conditions: Store at 4 °C. Avoid prolonged exposure to light.

**Application:** Flow Cytometry

**Background: CD9** belongs to proteins of tetraspanin family that orchestrate cholesterol-associated tetraspaninenriched signaling microdomains within the plasma membrane, forming complexes with each other as well as with integrins, membrane-anchored growth factors and other proteins. CD9 is involved in cell motility, osteoclastogenesis, neurite outgrowth, myotube formation, and sperm-egg fusion, plays roles in cell attachment and proliferation and is necessary for association of heterologous MHC II molecules on the dendritic cell plasma membrane which is important for effective T cell stimulation. CD9 is also considered as metastasis suppressor in solid tumors.

## References:

Schlossman S., L. Bloumsell, W. Gilks, et al., eds. 1995. Leucocyte Typing V: White Cell Differentiation Antigens. P: 246 Oxford University Press, New York.

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for <u>research use only</u>. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

## ImmunoTools Excellent Quality - Advantageously priced