

anti-human CD43 no azide

Monoclonal antibody MEM-59 to human CD43

Cat-No: **21270430**

100 µg in 100 µl

Clone: MEM-59

Specificity: The antibody MEM-59 recognizes neuraminidase-sensitive epitope on CD43 (Leukosialin), a 95-135 kDa type I transmembrane glycoprotein (mucin-type) which is involved in lymphocyte activation. CD43 is expressed by platelets and at high levels on the surface of all leukocytes; it is negative on resting B lymphocytes and erythrocytes.

HLDA IV; WS Code NL 604 / **HLDA V;** WS Code AS S290

Isotype subclass: Mouse IgG1

Form: Purified from cell culture supernatant by Protein-A affinity chromatography.

Purity: > 95% (by SDS-PAGE)

Physical state: Liquid

Buffer/Additives/Preservative: PBS (pH 7.2), sterile

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: Store at -20°C. Avoid freeze and thaw cycles.
Should be handled under aseptic conditions.

Application: Functional Application

References:

- 1.) Deckert, M. et al., Eur. J. Immunol. 22, 2943, 1992.
- 2.) Alvarado, M. et al., Eur. J. Immunol. 25, 1051, 1995.
- 3.) Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).
- 4.) Cermak, L. et al., J. Biol. Chem. 227, 7955, 2002.

Background: **CD43** (leukosialin, sialophorin) is a transmembrane mucin-like protein with high negative charge, expressed on the surface of most hematopoietic cells. CD43 contributes to a repulsive barrier that interferes with cellular adhesion, however, in certain cases also promotes leukocyte aggregation. By interaction with actin-binding proteins ezrin and moesin CD43 plays a regulatory role in remodeling T-cell morphology and regulates cell-cell interactions during lymphocyte traffic. CD43 signaling both enhances LFA-1 adhesiveness and counteracts LFA-1 induction via other receptors. Expression of CD43 causes induction of functionally active tumour suppressor p53 protein, but in case of p53 and ARF deficiency CD43 promotes tumour proliferation and viability. It appears to be an important modulator of leukocyte functions.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com