## anti-human CD44 no azide

## **Monoclonal Antibody MEM-85 to CD44 (Human)**

Cat-No: **21270440** 100 μg in 100 μl

Clone: MEM-85

**Specificity:** The antibody MEM-85 reacts with both cell surface-expressed and soluble form of CD44 antigen (Phagocyte glycoprotein 1), a 80-95 kDa transmembrane glycoprotein (hyaladherin family) present on the most of cells and tissues (leukocytes, endothelial cells, mesenchymal cells, etc.); it is negative on platelets and hepatocytes.

HLDA IV; WS Code NL 706 HLDA VI; WS Code AS Ref.15

Isotype subclass: Mouse IgG2b

Form: Purified from ascites by protein-A affinity chromatography.

**Purity:** > 95% (by SDS-PAGE)

Physical state: Liquid

Buffer/Additives/Preservative: sterile PBS, pH 7.2

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

Storage conditions: Store at 4 °C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.

**Application:** functional application

**Background: CD44** is a transmembrane glycoprotein expressed on the surface of most cells, which serves as a receptor for hyaluronan. CD44 mediates angiogenesis, cell adhesion, proliferation and migration, it is thus important for lymphocyte activation, recirculation and homing, it can thus serve e.g. as a modulator of macrophage recruitment in response to pathogen. Although CD44 functions are essential for physiological activities of normal cells, elevated CD44 expression correlates with poor prognosis in many carcinomas, facilitating tumour growth and metastasis, antiapoptosis and directional motility of cancer cells.

References: Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).

This material is offered for <u>research use only</u>. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.