

anti-rat CD161 NK-Cells (NKR-P1) no azide

Monoclonal Antibody 10-78 to CD161 (Rat)

Cat-No: **23151610**

100 µg in 100 µl

Clone: 10-78

Specificity: This anti-rat NK cell monoclonal antibody recognizes a 60 kDa disulfide linked homodimer known as NKR-P1. This type II transmembrane glycoprotein is found on rat NK cells and large granular lymphocytes. This antibody is suitable for flow cytometry. It is also reported to work with frozen sections.

Isotype subclass: Mouse IgG1

Form: purified

Physical state: Liquid

Buffer/Additives/Preservative: sterile PBS (pH 7.2).

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

Application: functional application

References:

- 1.) Ryan, J.C. et al (1991), NKR-P1, an activating molecule on rat natural killer cells, stimulates phosphoinositide turnover and a rise in intracellular calcium, *J Immunol.* 141, 3244-3250
- 2.) Sciba, A. et al. (1998), Phenotype of rat monocytes during acute kidney allograft rejection: increased expression of NKR-P1 and reduction of CD43. *Scand.J.Immunol.* 47, 332-342
- 3.) Josien, R. et al. (1997), Rat spleen dendritic cells express natural killer cell receptor protein 1 (NKR-P1) and have cytotoxic activity to select targets via Ca^{2+} -dependent mechanism, *J Exp. Med.* 186, 467-472
- 4.) Sciba, A. et al. (1997), Rat monocytes up-regulate NKR-P1A and down-modulate CD4 and CD43 during activation in vivo: monocyte subpopulations in normal and IFN- γ -treated rats. *J. Leuko.Bio.*, 62, 741-751
- 5.) Na, H. et al. (1992) Distribution of lymphocyte subsets in rat milk from normal and *Trichinella spiralis*-infected rats. *J.Repro.Bio.* 22, 269-279
- 6.) Giorda, R. et al. (1990) NKR-P1, a signal transduction molecule on natural killer cells, *Science*, 249, 1298-1300
- 7.) Vaage, J.T. et al. (1991), Allospecific recognition of hemiac cells in vitro by natural killer cells from athymic rats: evidence that allogen determinants coded for by single major histocompatibility complex haplotypes are recognized, *Eur.J.Immunol.* 21, 2167-2175
- 8.) Chambers, William H. et al. (1989), Monoclonal antibody to a triggering structure expressed on rat natural killer cells and adherent lymphokine-activated killer cells, *J Exp. Med.* 169, 1373-1389
- 9.) Chambers, W.H. et al. (1992) Functional heterogeneity between NKR-P1^{bright}/*Lycopersicon esculentum lectin* (L.E.)^{bright} and NKR-P1^{bright}/L.E.^{dim} subpopulations of rat natural killer cells, *J.Immunol.*, 148, 3658-3665 (no.11)

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com