

anti-mouse CD4 PE-conjugated

PE- conjugated monoclonal antibody GK1.5 to mouse CD4

Cat-No: **22850044**

500 µl

Clone: GK1.5

Specificity: The CD4 (L3/T4) antigen is thought to be the murine equivalent of the human Leu3/OKT4 antigen. This T cell surface molecule appears to be expressed by the helper/inducer subset of murine T cells and by delayed hypersensitivity T cells but not by cytotoxic T cells or their precursors. CD4 (L3/T4) and CD8a (Ly 2) have been shown to be present on mutually exclusive T cells in the peripheral lymphoid organs but the thymus contains cells expressing both CD4 (L3/T4) and CD8a (Ly2). The anti-mouse CD4 (L3/T4) mAb binds to approximately 85% of mouse thymocytes, 20% splenocytes, 50% lymph node cells, and a small number of bone marrow cells. It detects a protein of approximately 52 kDa on SDS-PAGE "Western Blots" (from Con A blast cell membranes) and is therefore similar to a well characterized human Leu3/T4 antigen (4).

Isotype subclass: Rat IgG2b

Form: The purified antibody is conjugated with R-Phycoerythrin (R-PE) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 1 % BSA and 0.09 % sodium azide (pH 7.2).

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: Store at 4 °C. Avoid prolonged exposure to light.

Application: Flow Cytometry

References:

- 1.) Teijaro, John R., David Verhoeven, Carly A. Page, Damian Turner, and Donna L. Farber. Journal of Virology 84, no. 18 (2010): 9217-9226.
2. Hafalla, Julius Clemence R., Carla Claser, Kevin N. Couper, Georges Emile Grau, Laurent Renia, J. Brian de Souza, and Eleanor M. Riley. PLoS Pathog 8, no. 2 (2012): e1002504.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com