

## anti-human/-porcine /-bovine CD21 PE-conjugated

PE-conjugated monoclonal Antibody LT21 to CD21

Cat-No: **21270214**

500 µl

**Clone:** LT21

**Specificity:** The antibody LT21 reacts with CD21 (CR2), a 145 kDa transmembrane glycoprotein (complement C3d receptor – C3dR) expressed on B lymphocytes, follicular dendritic cells, some epithelial cells and a subset of T lymphocytes. It is not expressed on immature B cells.

**HLDA VI; WS Code B CD21.1**

**Species Reactivity:** Human, Porcine, Bovine, Canine (Dog)

**Isotype subclass:** Mouse IgG1

**Form:** The purified antibody is conjugated with R-Phycoerythrin (R-PE) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing 1% BSA and 0.09% sodium azide (pH 7.2)

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

**Storage conditions:** Store at 4 °C. Avoid prolonged exposure to light

**Application:** Flow Cytometry

**Background:** CD21 (complement receptor 2, CR2) binds C3 complement fragments, especially its breakdown fragments, which remain covalently attached to complement activating surfaces or antigen. CD21 has important roles in uptake and retention of immunocomplexes, survival of memory B cells and in development and maintenance of the humoral response to T-dependent antigens. CD21 also serves as a key receptor for Epstein-Barr virus binding and is involved in targeting prions to follicular dendritic cells and expediting neuroinvasion following peripheral exposure to prions. A soluble form of the CD21 (sCD21) is shed from the lymphocyte surface and retains its ability to bind respective ligands.

**References:** Leukocyte Typing VI. Kishimoto T. et al. (Eds.), Garland Publishing Inc. (1997).

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for research only. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany  
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com  
www.immunotools.com