

## anti-human CD71 PE-conjugated

PE - conjugated monoclonal antibody MEM-75 to human CD71

Cat-No: **21270714**

500 µl

Clone: MEM-75

**Specificity:** The antibody MEM-75 reacts with CD71 antigen (transferrin receptor), a 95 kDa type II homodimeric transmembrane glycoprotein expressed on activated B and T lymphocytes, macrophages and erythroid precursors; it is lost on resting blood leukocytes.

The antibody MEM-75 does not block binding of transferrin to the receptor.

**HLDA IV; WS Code A 45**

**HLDA V; WS Code T T-165**

**Isotype subclass:** Mouse IgG1

**Form:** The purified antibody is conjugated with R-Phycoerythrin (R-PE) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing 1 % BSA and 0.09 % sodium azide (pH 7.2)

**Expiration date:** The reagent is stable until the expiry date stated on the vial label

**Storage conditions:** Store at 4 °C. Avoid prolonged exposure to light.

**Application:** Flow Cytometry

### References:

\*Rouault TA: 2003 Dec;1(3):E79.

\*Taketani S: 2005 Apr;205(4):297-318.


\*Graham Rmand other: 2007 Sep 21;13(35):4725-36.

\*Graham RM and other: 2008 Feb;48(2):327-34.

**Background: CD71** (transferrin receptor) is a type II transmembrane glycoprotein expressed as homodimer in erythroid blood cell line and in activated leukocytes. Upon binding of holotransferrin (complex of transferrin and iron ions), CD71 is internalized by clathrin-mediated endocytosis. Acidification of endosomes by vesicular membrane proton pumps leads to dissociation of iron ions, whereas transferrin (apotransferrin) remains associated with CD71 and recycles to the cell surface, where it is released upon exposure to normal pH. CD71 is also involved in uptake of non-transferrin bound iron.

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.



**Attention!** Cells from one healthy individual are shown. Cell Populations and staining intensity may vary interindividually.

**ImmunoTools** Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany

phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, [info@immunotools.com](mailto:info@immunotools.com)

[www.immunotools.com](http://www.immunotools.com)