

anti-human CD147 PE-conjugated

PE - conjugated monoclonal antibody MEM- M6/1 to human CD147

Cat-No: **21271474**

500 µl

Clone: MEM-M6/1

Specificity: The antibody MEM-M6/1 recognizes an epitope in the N-terminal Ig domain (D1) of CD147 (Neurothelin), a 50-60 kDa type I transmembrane glycoprotein primarily expressed on all leukocytes, red blood cells, platelets and endothelial cells; it is not expressed by resting lymphocytes. The antibody MEM-M6/1 is a high-affinity antibody capable of binding to unstimulated peripheral blood T cells.

Isotype subclass: Mouse IgG1

Form: The purified antibody is conjugated with R-Phycoerythrin (PE) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 1 % BSA and 0.09 % sodium azide (pH 7.2)

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: Store at 4 °C. Avoid prolonged exposure to light.

Application: The reagent is designed for Flow Cytometry analysis of human blood cells.

References:

- *Kirk P and others: EMBO J. 2000 Aug 1;19(15):3896-904.
- *Wilson MC and others: J Biol Chem. 2005 Jul 22;280(29):27213-21.
- *Xu D, Hemler ME: Mol Cell Proteomics. 2005 Aug;4(8):1061-71.
- *Iacono KT and others: Exp Mol Pathol. 2007 Dec;83(3):283-95.
- *Ruiz S and others: J Biol Chem. 2008 Feb 29;283(9):5554-66.

Background: **CD147** (basigin, neurothelin, OX-47, 5A11, CE9, M6) also known as EMMPRIN (extracellular matrix metalloproteinase inducer) or TCSF (tumour cell-derived collagenase-stimulatory factor) is an ubiquitously expressed cell surface protein with multiple glycosylated forms. The highest level of CD147 expression is on metabolically active cells, such as lymphoblasts, inflammatory cells, brown adipocytes and malignant tumour cells. CD147 has multiple functions, including facilitating of cell surface expression of monocarboxylate transporter proteins and extracellular matrix metalloproteinases, regulation of integrin functions, it plays roles in cell development and activation, fetal development or retinal function.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com