

anti-human CD69 PE-conjugated

PE- conjugated monoclonal Antibody FN50 to CD69 (Human)

Cat-No: **21620694**

500 µl

Clone: FN50

Specificity: The antibody FN50 recognizes the CD69 antigen, an early activation marker (within 2h) on T, B and NK-cells, possibly involved in signal transduction. It is also expressed by activated platelets. On flow cytometry the antibody stains > 90% of activated human peripheral blood lymphocytes. This antibody has been studied at the 4. International Workshop on Human Leucocyte Differentiation Antigens.

Isotype subclass: Mouse IgG1/kappa

Form: The purified antibody is conjugated with R-Phycoerythrin (R-PE) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 1% BSA and 0.09% sodium azide (pH 7.2)

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: Store at 4 °C. Avoid prolonged exposure to light.

Application: The FN50 antibody is well suited for detection of CD69 in flow cytometry.


References:

- *López-Cabrera M and other: 1993 Aug 1;178(2):537-47.
- *Nielsen SD and other: 1998 Oct;114(1):66-72.
- *Pitsios C and other: 2008;68(3):233-41.
- *Konjević G and other: 2007 Nov;37(11):887-96.

Background: CD69 (C-type lectin domain family 2 C, CLEC2C, also known as AIM) is one of the earliest inducible cell surface molecules acquired during leukocyte activation. This glycoprotein serves as a lectin-type receptor in lymphocytes, NK cells and platelets; it is involved in lymphocyte proliferation. CD69 expression is counteracted on T cells in the AIDS stage of HIV infection, and may be also predictive for clinical response to chemoimmunotherapy.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.



Attention! Cells from one healthy individual are shown. Cell Populations and staining intensity may vary interindividually.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany

phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com

www.immunotools.com