

## anti-human CD56 PE-conjugated

PE - conjugated monoclonal antibody B-A19 to human CD56

Cat-No: **21810564**

500 µl

**Clone:** B-A19

**Specificity:** The antibody B-A19 reacts with a 175-220 kDa isoform of CD56 (NCAM) expressed in leukocytes. It has been suggested that the antibody B-A19 could react with rhesus monkey lymphocytes. Reactivity with other NCAM isoforms has not been tested.

**HLDA VI; WS code A055 - HLDA VI; WS Code NK26 - HLDA VII; WS code 70077**

**Isotype subclass:** Mouse IgG1

**Form:** The purified antibody is conjugated with R-Phycoerythrin (R-PE) under optimum conditions. The reagent is adjusted for direct use. No reconstitution is necessary.

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing 1 % BSA and 0.09 % sodium azide (pH 7.2)

**Expiration date:** The reagent is stable until the expiry date stated on the vial label

**Storage conditions:** Store at 4 °C. Avoid prolonged exposure to light.

**Application:** Flow Cytometry

### References:

\*Jakovcevski I, Mo Z, Zecevic N: Neuroscience. 2007 Oct 26;149(2):328-37.

\*Ohishi Y, Kaku T, Oya M, et al: Gynecol Oncol. 2007 Oct;107(1):30-8.

\*McCluggage WG, McKenna M, McBride HA: Int J Gynecol Pathol. 2007 Jul;26(3):322-7.

**Background:** **CD56** (NCAM, neural cell adhesion molecule) is a transmembrane glycoprotein of immunoglobulin family serving as adhesive molecule which is ubiquitously expressed in nervous system, usually as 120 kDa, 140 kDa or 180 kDa isoform, and it is also found on T cells and NK cells. Polysialic modification results in reduction of CD56-mediated cell adhesion and is involved in cell migration, axonal growth, pathfinding and synaptic plasticity. CD56 is a widely used neuroendocrine marker with a high sensitivity for neuroendocrine tumours and ovarian granulosa cell tumours.

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany  
phone: +49-(0)4491-400997, fax: +49-(0)4491-400998, info@immunotools.com  
www.immunotools.com