anti-human/-rat/-bovine CD54 PE/Dy647-conjugated

PE/Dy647 -conjugated monoclonal antibody 1H4 to human CD54

Cat-No: **21279547** 500 μI

Clone: 1H4

Specificity: The antibody 1H4 reacts with CD54 (ICAM-1), a 85-110 kDa type I transmembrane glycoprotein (receptor for rhinovirus) expressed on activated endothelial cells, T lymphocytes, B lymphocytes, macrophages, granulocytes and dendritic cells; the expression of CD54 is upregulated by activation.

Isotype subclass: Mouse IgG2b

Species Reactiity: human, rat, bovine

Form: The purified antibody is conjugated with PE-Dyomics 647 (PE-Dy647) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No reconstitution is necessary.

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 1% BSA and 0.09% sodium azide (pH 7.2)

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: Store at 4 °C. Do not freeze. Avoid prolonged exposure to light.

Application: The reagent is designed for Flow Cytometry analysis.

References:

- *Boyd AW and others: 1988 May;85(9):3095-9.
- *Boyd AW and othes: 1989 May 15;73(7):1896-903.
- *Springer TA.: 1990 Aug 2;346(6283):425-34.
- *Ockenhouse CF and othes: 1992 Jan 10;68(1):63-9. Erratum in: Cell 1992 Mar 6;68(5):following 994.

Background: CD54 (ICAM-1) is a 90 kD member of the C2 subset of immunoglobulin superfamily. It is a transmembrane molecule with 7 potential N-glycosylated sites, expressed on resting monocytes and endothelial cells and can be upregulated on many other cells, e.g. with lymphokines, on B- and T-lymphocytes, thymocytes, dendritic cells and also on keratinocytes, chondrocytes, as well as epithelial cells. CD54 mediates cell adhesion by binding to integrins CD11a/CD18 (LFA-1) and to CD11b/CD18 (Mac-1). The interaction of CD54 with LFA-1 enhances antigen-specific T-cell activation.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for <u>research use only</u>. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.