

anti-rat CD8b

Monoclonal Antibody 3.4.1. to CD8b (Rat)

Cat-No: **23159081**

100 µg in 100 µl

Clone: 3.4.1

Specificity: This anti-rat CD8b monoclonal antibody reacts with beta chain of the CD8 differentiation antigen. CD8b is expressed on most thymocytes and mature T cytotoxic/suppressor cells (MHC class I restricted). While the CD8a and CD8b form a heterodimer on the surface of thymocytes and thymus dependent T cytotoxic/suppressor cells, the majority of NK cells, many CD8 T cells from athymic rats, many activated CD4 T cells, and intestinal epithelium lymphocytes (IEL) express CD8a without CD8b. This suggests that expression of the CD8 heterodimer (a/b) is more dependant on intrathymic T cell maturation than that of the homodimer (a/a). The thymus dependence of CD8a/b T cells may be due to a requirement for thymic selection on self MHC class I antigens. Reported applications for this antibody include flow cytometry, immunoprecipitation and Western blotting. The 3.4.1 antibody also blocks both activation in an allogenic response and cell mediated cytotoxicity by CD8 T cells.

Isotype subclass: Mouse IgG1K

Form: purified

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 1 % BSA and 0.09 % sodium azide (pH 7.2).

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: For long term storage, aliquot and freeze unused portion at -20°C in volumes appropriate for single usage. Avoid freeze/thaw cycles.

Application: Flow Cytometry, Immunoprecipitation and Western blotting

References: 1. Elflein, K., Rodriguez-Palmero, M., et al. (2003) Immunobiology. 102, 1764-1770.
2. Torres-Nagel, N, Kraus, E., Brown, M.H., Tiefenthaler, G., Mintnacht, R., Williams, A.F., and T. Hunig. (1992) Eur. J. Immunol. 22, 2841-2848.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink, and animal feedingstuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Altenoyther Str. 10; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com