

# anti-human CD40

## Monoclonal Antibody HI40a to CD40 (Human)

Cat-No: **21270401**

100 µg in 100 µl

**Clone:** HI40a

**Specificity:** The antibody HI40a recognizes CD40 (BP50), a 48 kDa type I single chain transmembrane glycoprotein expressed on normal and neoplastic B cells, but not on terminally differentiated plasma cells. CD40 antigen is also present on Hodgkin's and Reed-Sternberg cells, follicular dendritic cells, some macrophages, basal epithelial cells and endothelial cells.

**Isotype subclass:** Mouse IgG1

**Form:** Purified from ascites by protein-G affinity chromatography.

**Purity:** > 95% (by SDS-PAGE)

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing 0.09 % sodium azide (pH 7.2).

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

**Storage conditions:** Store at 4°C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.

**Application:** Flow Cytometry

### References:

- \*) Oxenius A, Campbell KA, Maliszewski CR, Kishimoto T, Kikutani H, Hengartner H, Zinkernagel RM, Bachmann MF. J Exp Med. 1996 May 1;183(5):2209-18
- \*) Grewal IS, Flavell RA: Immunol Rev. 1996 Oct;153:85-106.
- \*) Pearson LL, Castle BE, Kehry MR: Int Immunol. 2001 Mar;13(3):273-83.

**Background:** CD40 is a costimulatory molecule of the TNF receptor superfamily and is expressed on many cell types, such as B cells, monocytes/macrophages, dendritic cells, endothelial cells, fibroblasts or vascular smooth muscle cells. Interaction of CD40 and its ligand CD154 (CD40L) is required for the generation of antibody responses to T-dependent antigens as well as for the development of germinal centers and memory B cells. In monocytes/macrophages CD40 engagement induces production of pro-inflammatory cytokines and chemokines. CD40-CD154 interactions are also critical for development of CD4 T cell-dependent effector functions. CD40 links innate and adaptive immune responses to bacterial stimuli and serves as an important regulator affecting functions of other costimulatory molecules.

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany  
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com  
www.immunotools.com