

anti-human CD63

Monoclonal Antibody MEM-259 to CD63 (Human)

Cat-No: **21270631**

100 µg in 100 µl

Clone: MEM-259

Specificity:

The antibody MEM-259 reacts with CD63 (LAMP-3), a 40-60 kDa tetraspan glycoprotein expressed by granulocytes, platelets, T cells, monocytes/macrophages and endothelial cells. Cell surface exposition of CD63 is usually activation-dependent.

Isotype subclass: Mouse IgG1

Form: Purified from ascites by protein-A affinity chromatography.

Purity: > 98% (by SDS-PAGE)

Physical state: Liquid

Buffer/Additives/Preservative: PBS with 0.1% sodium azide (pH 7.2).

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: Store at 4°C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.

Application: Flow Cytometry: detection of activated platelets, neutrophils and basophils
Immunoprecipitation
Western Blotting: non reducing conditions

References:

- *Grützkau A and other: 2004 Sep;61(1):62-8.
- *Mantegazza AR and other: 2004 Aug 15;104(4):1183-90.
- *Pfistershammer K and other: 2004 Nov 15;173(10):6000-8.
- *Israels SJ, McMillan-Ward EM: 2005 Feb;93(2):311-8.
- *Kwon MS and other: 2007 Jul;57(1):46-53.
- *Lin D and other: 2008 Jan 22

Background: **CD63** (LAMP-3, lysosome-associated membrane protein-3), a glycoprotein of tetraspanin family, is present in late endosomes, lysosomes and secretory vesicles of various cell types. It is also present in the plasma membrane, usually following cell activation. Hence, it has become an widely used basophil activation marker. In mast cells, however, CD63 exposition does not need their activation. CD63 interacts with integrins and affects phagocytosis and cell migration, it is also involved in H/K-ATPase trafficking regulation of ROMK1 channels. CD63 also serves as a T-cell costimulation molecule. Expression of CD63 can be used for predicting the prognosis in earlier stages of carcinomas.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com