

# Mouse IgG1 Isotype control

## monoclonal antibody MOPC-21 to IgG1

Cat-No: **21275611**

100 µg in 100 µl

**Clone:** MOPC-21

**Specificity:** This mouse IgG1 monoclonal antibody (clone MOPC-21) has unknown specificity and was chosen as an isotype control after screening on variety of resting, activated, live and fixed rat and human tissues.

**Negative Species:** human, rat.

**Isotype subclass:** Mouse IgG1

**Purity:** > 98% (by SDS-PAGE)

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS containing 0.09% sodium azide, (pH 7.2)

**Storage conditions:** Aliquot and store at -20°C. Avoid freeze/thaw cycles.

**Application:** Western Blots, Immunohistochemistry( p/f), Immunoprecipitation, Flow Cytometry, Control Experiments

**Background:** The specificity of staining by monoclonal antibodies to target antigens should be verified by establishing the amount of non-specific antibody binding. Especially at higher concentration (more than 15 µg/ml) the antibody staining usually has consignable background. To this end a non-reactive immunoglobulin of the same isotype is included as a negative control for each specific monoclonal antibody used in a particular immunoassay. The monoclonal antibody MOPC-21, generated against an undefined antigen, does not react specifically with rat and human samples, and hence all the background that could be observed when working with this antibody would be a result of general nonspecific interactions between an mouse IgG1 molecule and the respective sample under the particular conditions. This shall help the customer to set up the experimental conditions so that the nonspecific binding of any antibody is abolished.

Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research only**. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Altenoyther Str. 10; 26169 Friesoythe; Germany  
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, [info@immunotools.com](mailto:info@immunotools.com)  
[www.immunotools.com](http://www.immunotools.com)