

# anti-human p53

## Monoclonal Antibody BP53-12 to p53 (Human)

Cat-No: **21279961**

100 µg in 100 µl

**Clone:** BP53-12

**Specificity:** The antibody BP53-12 recognizes defined epitope (aa 16-25) on human p53, a 50 kDa tumour suppressor found in increased amounts in a wide variety of transformed cells; it is frequently mutated or inactivated in many types of cancer.

**Isotype subclass:** Mouse IgG2a

**Form:** Purified from ascites by protein-A affinity chromatography.

**Purity:** > 95% (by SDS-PAGE)

**Physical state:** Liquid

**Buffer/Additives/Preservative:** PBS with 0.09 % sodium azide (pH 7.2).

**Expiration date:** The reagent is stable until the expiry date stated on the vial label.

**Storage conditions:** Store at 4 °C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles.

**Application:** Immunocytochemistry, Immunohistochemistry (p), Western Blotting, ELISA, Immunoprecipitation

### References:

- \*Agarwal ML and others: Proc Natl Acad Sci U S A. 1998 Dec 8;95(25):14775-80.
- \*Taylor WR and others: Mol Biol Cell. 1999 Nov;10(11):3607-22.
- \*Taylor WR and others: Oncogene. 1999 Jan 14;18(2):283-95.
- \*Tanigawa S and others: Biosci Biotechnol Biochem. 2008 Mar;72

**Background:** The tumour suppressor protein p53 is a key element of intracellular anticancer protection. It mediates cell cycle arrest or apoptosis in response to DNA damage or to starvation for pyrimidine nucleotides. It is up-regulated in response to these stress signals and stimulated to activate transcription of specific genes, resulting in expression of p21waf1 and other proteins involved in G1 or G2/M arrest, or proteins that trigger apoptosis, such as Bcl-2. The structure of p53 comprises N-terminal transactivation domain, central DNA-binding domain, oligomerisation domain, and C-terminal regulatory domain. There are various phosphorylation sites on p53, of which the phosphorylation at Ser15 is important for p53 activation and stabilization.

**Warning:** Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for research only. Not for use in human. For in vitro use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

**ImmunoTools** Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany  
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, [info@immunotools.com](mailto:info@immunotools.com)  
[www.immunotools.com](http://www.immunotools.com)