

anti-human CD66eb (CEACAM5/8)

Monoclonal antibody 5C8C4 to human CD66eb (CEACAM5/8)

Cat-No: 21608661

100 µg in 100 µl

Clone: 5C8C4

Specificity: The antibody 5C8C4 recognizes the CEACAM5 (CEA) antigen, a 180 kDa GPI-anchored glycoprotein expressed on various human epithelial cells and the CEACAM8 antigen, a 95 kDa GPI-anchored glycoprotein expressed solely on human granulocytes.

Isotype subclass: Mouse IgG1 kappa.

Form: purified

Physical state: Liquid

Buffer/Additives/Preservative: PBS containing 0.09 % sodium azide (pH 7.2)

Expiration date: The reagent is stable until the expiry date stated on the vial label.

Storage conditions: Store at 4 °C. For long-term storage aliquot and store at -20°C. Avoid freeze/thaw cycles

Application: Flow Cytometry, ELISA, IP, WB, IHC (f/p), ICC

Background: **CD66e**, also known as CEACAM5 (Carcinoembryonic antigen-related (CEA) cell adhesion molecule 5, is a glycosyl phosphatidyl inositol- (GPI-) anchored glycoprotein. CEACAM5 and related genes make up the CEA family belonging to the immunoglobulin superfamily. CD66e is involved in cell adhesion. Additionally it serves as pathogen receptor (e.g. Helicobacter pylori, Neisseria, Moraxella, Candida albicans). Antibodies against CEACAM5 are commonly used in immunohistochemistry to identify cells expressing the glycoprotein in tissue samples. However, CEACAM5 is also found in serum where it can be used as a tumor marker. In adults, CEACAM5 is primarily expressed in epithelia of the gastrointestinal tissue commonly together with CEACAM1, CEACAM6 and CEACAM7. **CD66b**, also known as CEACAM8 is a GPI-anchored glycoprotein as well. CEACAM8 is involved cell-cell communication* but does NOT serve as pathogen receptor. In adults CEACAM8 is expressed in granulocytes together with CEACAM1, CEACAM3 and CEACAM6.

References:

- 1.) Singer BB, Scheffrahn I, Kammerer R, Suttrop N, Ergun S, Slevogt H: Deregulation of the CEACAM expression pattern causes undifferentiated cell growth in human lung adenocarcinoma cells. PLoS One. 2010 Jan 18;5(1):e8747.
- 2.) Gangopadhyay A, Bajenova O, Kelly TM, Thomas P: Carcinoembryonic antigen induces cytokine expression in Kupffer cells: implications for hepatic metastasis from colorectal cancer. Cancer Res. 1996 Oct 15;56(20):4805-10.

Warning: Sodium azide is harmful if swallowed (R22). Keep out of reach of children (S2). Keep away from food, drink and animal feeding stuff (S13). Wear suitable protective clothing (S36). If swallowed, seek medical advice immediately and show this container or label (S46). Contact with acids liberates very toxic gas (R32). Azide compounds should be flushed with large volumes of water during disposal to avoid deposits in lead or copper plumbing where explosive conditions can develop.

This material is offered for **research use only**. Not for use in human. For *in vitro* use only. ImmunoTools will not be held responsible for patent infringement or other violations that may occur with the use of our products.

ImmunoTools Excellent Quality - Advantageously priced

Gladiolenweg 2; 26169 Friesoythe; Germany
phone:+49-(0)4491-400997, fax:+49-(0)4491-400998, info@immunotools.com
www.immunotools.com